

U. PORTO

FMUP FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2017/2018

Maria Clotilde Avides Moreira Pinto Osório

**Uma visão geral no tratamento da
síndrome hemolítica urémica atípica**

**An overview on the treatment for
atypical hemolytic uremic syndrome**

março, 2018

FMUP

Maria Clotilde Avides Moreira Pinto Osório
Uma visão geral no tratamento da
síndrome hemolítica urémica atípica
An overview on the treatment for atypical
hemolytic uremic syndrome

Mestrado Integrado em Medicina

Área: Ciências médicas e da saúde

Tipologia: Monografia

Trabalho efetuado sob a Orientação de:

Dr.^a Ines Passos e Castro Neto Ferreira

Trabalho organizado de acordo com as normas da revista:

Seminars in Nephrology

março, 2018

FMUP

Eu, Maria Clotilde Avides Moreira Pinto Osório, abaixo assinado, nº mecanográfico 201104867, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 17/03/2018

Assinatura conforme cartão de identificação:

Maria Clotilde Osório

NOME

Maria Clotilde Avides Moreira Pinto Osório

NÚMERO DE ESTUDANTE

E-MAIL

201104867 mclotilde.p@ gmail . com

DESIGNAÇÃO DA ÁREA DO PROJECTO

Ciências médicas e da saúde

TÍTULO DISSERTAÇÃO/MONOGRAFIA (riscar o que não interessa)

An overview on the treatment for atypical hemolytic uremic syndrome

ORIENTADOR

Dr.ª Inês Passes e Castro Neto Ferreira

COORDENADOR (se aplicável)

—

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTES TRABALHOS APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input checked="" type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTES TRABALHOS (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTES TRABALHOS.	<input type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 17/03/2018

Assinatura conforme cartão de identificação: Maria Clotilde Osório

Aos meus avós Teresa e António, por me transmitirem a paixão pela medicina e pelas pessoas. Aos meus pais, ao Nicolau, à Maria e ao Augusto por todos os dias me mostrarem o verdadeiro significado de família. Ao Filipe pelo agora, pelo o que já passou, pelo que virá.

An overview on the treatment for atypical hemolytic uremic syndrome

Maria Clotilde Avides Moreira Pinto Osório^{a)}, Inês Passos e Castro Neto Ferreira^{b)}

^{a)} Medical Student from Faculdade de Medicina da Universidade do Porto

^{b)} MD, Nephrology Department, Centro Hospitalar São João Porto; Instituto de Inovação e Investigação em Saúde (I3S); Grupo de I&D em Nefrologia e Doenças Infecciosas; Instituto Nacional de Engenharia Biomédica (INEB); Faculdade de Medicina da Universidade do Porto, Porto

From Faculdade de Medicina da Universidade do Porto, Alameda Prof. Hernâni Monteiro, 4200-319 Porto, Portugal

Contact: +351967168406/ mclotilde.po@gmail.com / Estrada do Marinho 8, 4970-630 Arcos de Valdevez, Portugal

ABSTRACT

Atypical hemolytic uremic syndrome (aHUS) is characterized by microangiopathic hemolytic anemia, thrombocytopenia, and acute renal failure caused by a dysregulation in the complement system, specifically in the alternative pathway. These abnormalities are due to a genetic mutation in half of the cases, and the response to the treatment and the prognosis vary accordingly to which regulatory mechanism is impaired. Endothelial cell swelling and detachment, arterial wall thickening, namely glomerular capillary wall and mesangiolysis are the main renal histologic findings. The first step in patient approach will be the exclusion of other thrombotic microangiopathies such as typical hemolytic uremic syndrome (STEC-HUS), thrombotic thrombocytopenic purpura (TTP) among others since the clinical presentation is very similar. The treatment of aHUS consists in supportive measures namely plasma exchange, which should be started 24 hours of the patient's presentation. Eculizumab, a recombinant humanized monoclonal antibody, is the only agent approved for the treatment of non-Shiga toxin by FDA, which effectiveness has been reported. In those studies, eculizumab led to improvement of hematologic manifestations and renal function, cessation of plasma therapy, including in several cases that had not responded to plasma exchange. Renal transplantation is an option for patients that develop end-stage renal disease, albeit the high recurrence rates, and hepatic transplantation is reserved for special cases.

The high mortality and morbidity contributes for the poor prognosis of these group of patients. This article will approach the general characteristics of aHUS, its treatment

options, the correct management and monitoring of the disease response, the outcomes and the proposed recent therapies.

KEYWORDS: Atypical hemolytic uremic syndrome; complement system; plasma therapy; eculizumab; renal transplantation.

INTRODUCTION

Hemolytic uremic syndrome (HUS) belongs to the large and diverse group of the thrombotic microangiopathies (TMAs) and is a life-threatening disorder characterized by a microangiopathic hemolytic anemia (MAHA), nonimmune, thrombocytopenia, and acute renal failure.^(1, 2) We can consider two types of HUS. The most common (with a frequency of more than 90%) is classified as typical HUS or Shiga toxin-producing *Escherichia coli* HUS (STEC-HUS) and is due to a bacterial infection producing a Shiga-like toxin (Stx1 and Stx2), most frequently due to O157:H7 serotype enterohemorrhagic *Escherichia coli* infection.⁽³⁾ It is more frequent in pediatric age, and often results in diarrhea, with favorable prognosis since as many as 70-85% of patients recover renal function.⁽⁴⁾

The others 10% of all cases of the HUS are categorized as atypical HUS (aHUS), a rare disorder caused by a dysregulation of the complement alternative pathway due to an impairing of the regulatory mechanisms. The outcome is poor due to the high risk of progression to end-stage renal disease (ESRD) and death.⁽⁵⁾

More than 50% of the aHUS cases have genomic defects and/or autoantibodies against complement regulatory proteins.^(6, 7) The penetrance of the disease is reported as incomplete in all predisposing genes, highlighting the requirement of a precipitating event like drugs, autoimmune conditions, cancer, transplants, pregnancy, or metabolic conditions, for disease manifestation.⁽⁸⁾

When it comes to treating a patient with aHUS treatment should be applied as soon as possible. Plasma therapy (PT) remains the initial treatment, preferably with intensive plasma exchange (PE) at high volumes and with a variable frequency, based on disease activity.⁽⁹⁾ The clinical response after 5-7 days of daily PE treatment should aid further decisions since the complete hematological and renal recovery rates are usually below 50% and differ according to the type of mutations or the levels of antibody titers.^(5, 9)

The discovery that complement system played an important role in aHUS led to the development of complement-inhibiting therapy.⁽⁹⁾ Eculizumab, a humanized monoclonal antibody against C5, blocks the formation of the membrane attack complex (MAC), stopping the hemolytic process, allowing the interruption of PT for long-term, with improvements in renal function and in the prognosis.⁽⁶⁾

Other options like renal and/or liver transplantation, could also be considered. New therapeutic strategies are also being studied.^(1, 2)

EPIDEMIOLOGY

The information available about the epidemiology of aHUS is limited, but a multicenter study in Europe stated an aHUS incidence of 0.11 cases/million habitants and according to the European Medicines Agency (EMA) the prevalence may be of 3.3 patients/million habitants/year among patients <18 years old, corresponding to 40 to 60% of all aHUS cases. Male and female are equally affected but the prevalence rises among women when the disease develops in adulthood.^(6, 10)

PATHOGENESIS

The complement system is part of the innate system but it also participates in the adaptive immunity. It's activated by the classical, the lectin or by the alternative pathway.⁽⁸⁾ The last one is constantly low-activated in plasma, and regulation defects or

over-activation ones, lead to cellular attack amplification, culminating in cell apoptosis and tissue damage, as seen in aHUS.^(2, 8) This complex mechanism has also multiple amplification loops which help maintaining the disease process, as seen on Fig. 1.

The regulation of the complement system is performed by membrane proteins such as *factor H*, *CD35 (complement receptor type 1)*, *CD46 (MCP)*, *CD55 (decay accelerating factor)* that act as cofactors for *plasma protein factor I* leading to the cleavage of *C3b* to inactive *C3b* and dissociating *C3* and *C5 convertases*. This regulation is also accomplished by *CD59 (MAC-inhibitory protein)*, another membrane protein, which promotes the cessation of the lytic membrane attack complex.^(2, 11)

Genetic mutations have been described in 50 to 60% of aHUS, comprising both familial and sporadic forms.^(2, 6)

Familial cases represent less than 20% of aHUS patients with both dominant and recessive hereditary forms described. Autosomal recessive aHUS often occurs in childhood. On the other hand, autosomal dominant aHUS affects mainly adults.⁽¹²⁾

Nearly 50% of the sporadic forms appear to be idiopathic.⁽²⁾

The activity of the complement regulatory proteins can be decreased by loss-of-function mutations (*CFH*, *CFI*, *CD46*, and *THBD*) or due to autoantibodies against *complement factor I* or *factor H*. The occurrence of these antibodies has an association with the homozygous *CFHR1-3* deletion, showing that they aren't strictly acquired. Conversely, the increase of the complement alternative pathway activation can also occur by gain-of-function mutations in *CFB* or *C3*.⁽⁵⁾

Mutations in a non-complement protein, including *diacylglycerol kinase ϵ* , *plasminogen* and *factor XII*, have also been recognized, and in some cases in association to *factor H autoantibodies*.^(11, 13)

More than 3% of the patients with aHUS have combined mutations.⁽⁶⁾

The penetrance of the disease is incomplete since only 50% of the mutations carriers manifest aHUS. However, the percentage can increase in patients with combined mutations.^(8, 10)

Patients who are carriers of complement mutations may stay asymptomatic for decades, suggesting that a trigger seem be essential for disease manifestation.^(1, 2) aHUS can be triggered by infections, pregnancy or post-partum period, autoimmune diseases, cancer, bone marrow or solid organ transplantation, and drugs such as immunotherapeutic agents (calcineurin inhibitors or the mammalian target of rapamycin inhibitors (mTORi)), platelet antiaggregants (e.g., ticlopidine and clopidogrel), antivirals, oral contraceptive drugs and chemotherapeutic drugs.^(8, 14, 15) Infections are responsible for at least half of these cases, mostly upper respiratory tract infections (influenza H1N1 virus) or non-*E. coli* diarrhea/gastroenteritis.^(6, 8) Pregnancy, mostly during post-partum period, is a frequent trigger of aHUS among women.⁽¹⁶⁾ The prevalence, mechanisms, response to treatment and prognosis differ to each mutation, as shown in Tab.1.

The hemolysis and the formation of microthrombi in aHUS can be explained by the effects that complement system has on platelets, red cells and endothelial cells. The complement activation can be induced by endothelial cell damage, promoting damage-induced enhancement of complementary activation and culminating in further damage of nearby cells as shown on Fig.2. Although this activation is more probable to happen via the alternative pathway in aHUS, classical and lectin pathways can also participate because of the exposure of novel epitopes recognized by both. Nonetheless, it was not identified yet which cells are damage first, being as potential candidates: the complement-mediated cell damage, direct activation of platelets, pro-coagulative state

in the endothelium, dysregulation of the coagulation or thrombolytic pathways, and direct complement-independent cell damage.⁽¹¹⁾

CLINICAL MANIFESTATIONS

The consequence of thrombosis and ischemic injury of multiple organs is seen on aHUS clinical manifestations.⁽¹⁾ Symptoms are consistent with severe MAHA, thrombocytopenia and acute renal failure such as pallor, fatigue, decreased urine output and edema. Hypertension, which results of hyperreninemia and volume expansion, can be a new onset or an exacerbation of a previously known disease.⁽¹⁰⁾ 20% of the patients can present extra-renal manifestations, such as neurological symptoms (like altered conscious, focal neurologic deficits, seizures, encephalopathy, hemiparesis, visual abnormalities, hemiplegia and coma), cardiovascular manifestations (myocardial infarction, cardiomyopathy, heart failure, and peripheral ischaemic vasculopathy) and gastrointestinal signs (prodromic diarrhea, colitis, nausea, vomit, abdominal pain, hepatitis, cholestasis and pancreatitis).^(5, 6, 8) Even though the presentation is often abrupt, aHUS progression can be insidious in 20% of the cases.⁽¹⁾

Laboratory findings include anemia, high levels of lactate dehydrogenase (LDH), reduced haptoglobin levels and the presence of schistocytes on a peripheral blood smear (all related to MAHA), thrombocytopenia, elevated serum creatinine level, mild proteinuria and microscopic hematuria.^(1, 5, 6) Prothrombin and partial thromboplastin times are normal, serum *C4* concentration is normal and *C3* levels may be reduced.^(1, 10)

aHUS is impossible to distinguish from typical HUS on standard histological analysis.⁽²⁾

In the acute phase endothelial cell swelling and detachment frequently come along with thrombosis, although thrombi are not essential to a histological diagnosis of aHUS.⁽¹⁾

Fibrinoid necrosis can be found in glomerular capillaries and arterioles, consisting of fibrin, cellular debris and rare neutrophils.⁽¹⁾ Glomerular capillary and arterioles wall thickening are often observed because the integrity of the vascular endothelial cells is lost, leading to accumulation of plasma proteins in the subendothelial zone. If a comparable material accumulates in the mesangium, mesangiolysis and glomerular capillary aneurysms can occur.⁽¹⁷⁾ Finally, mucoid intimal hyperplasia narrows the vascular lumen.⁽⁸⁾ Immunohistological examination shows erratic deposition of fibrin, IgM, C3 and C1q in affected areas.⁽¹⁾

DIAGNOSIS AND DIFFERENTIAL DIAGNOSIS

The first step of the diagnosis approach of a thrombotic microangiopathy patient is to distinguish between STEC-HUS, aHUS and thrombotic thrombocytopenic purpura (TTP).⁽⁸⁾ A complete clinical history, including family medical history, physical examination and hematological, biochemical and microbiological assays should be performed.⁽¹⁷⁾ However, all blood samples must be executed before PT.⁽⁶⁾

When comparing STEC-HUS with atypical HUS there are some similarities to consider, such as the clinical signs and symptoms.^(10, 11) Furthermore, the pathogenesis is identical, since endothelial cells, red cells and platelets are affected in both entities.⁽¹¹⁾

A positive *Shiga toxin* teste in stool samples and a positive Sorbitol-MacConkey stool culture (detecting E coli O157:H7, the most common STEC) validate the diagnosis of STEC-HUS.⁽³⁾ A urine sample can also be undertaken to exclude a urinary STEC infection.

The Coombs test must be also undertaken, and in case of a positive result, the possibility of pneumococcal HUS must be considered.⁽¹⁰⁾

Establishing the diagnosis of TTP or HUS based only in clinical criteria isn't reliable because of the similarity of both presentations.⁽¹⁵⁾ Therefore to exclude TTP it's mandatory to confirm a normal ADAMTS13 (a disintegrin and metalloproteinase with thrombospondin type 1 motif 13) activity (>5-10%).^(6, 10)

Scleroderma renal crisis, antiphospholipid syndrome, malignant hypertension, malignancy-related TMA, methylmalonic aciduria, homocystinuria type C disseminated intravascular coagulation (DIC) and HELLP (hemolysis, elevated liver enzymes, and thrombocytopenia) syndrome are other forms of TMA that must be excluded.^(1, 5)

The diagnosis of aHUS is a diagnostic of exclusion.⁽⁶⁾ The hematological and histological findings referred above are important when suspicion of aHUS diagnosis is made, albeit in some cases, schistocytes could be absent and platelet count or serum C3 concentration may be normal.^(1, 6) Elevated serum C3 concentration is present in 30 to 50% of aHUS cases but could also point to a C3 glomerulopathy (C3G), highlighting the importance of renal biopsy for diagnosis. Contrary to aHUS, C3G follows a chronic and indolent progression with persistent alternative pathway activation, resulting in predominant glomerular C3 fragment deposition and a 10-years renal survival rate of nearly 50%.⁽⁷⁾

Complement analysis requires serum levels of *C3*, *C4*, *CFH*, and *CFI*.⁽⁸⁾ In aHUS patients is expected a normal CH50 value (antibody-coated sheep erythrocyte lysis assay that determines the total complement activity) and a reduced AP50 level (a rabbit erythrocyte lysis assay that determines the alternative complement activity). Individuals with *FH* mutation or *FH* autoantibodies present an abnormal *CFH* functional assay with

sheep erythrocytes lysis. A new functional assay called modified Ham test has been recently suggested as promising test for the distinction of aHUS from STEC-HUS, TTP and DIC, highlighting future improvement in the diagnostic of TMA.⁽¹⁰⁾

Screening for *MCP* mutations can be made by a fast and an inexpensive method, such as by flow cytometry on mononuclear cells, although genetic analysis is still required to detect all changes.⁽⁸⁾

Nowadays, the diagnosis of aHUS is based on genetic studies, even though the proportion of variants identified are still a challenge due to the uncertainty of its clinical significance. Further functional assays are needed to evaluate its relevance.^(8, 10) Despite that, genetic screening in these patients has also diagnostic and prognostic relevance and might influence the natural course and the long-term management of the disease.⁽¹⁰⁾

Renal biopsy is not indicated in patients with a solid diagnosis of aHUS (e.g. positive family history or recurrence) due to the risk of bleeding. However, it may be needed for acute renal failure in adult patients to verify the etiology, exclude other processes and assess the prognosis.⁽⁶⁾

TREATMENT

A thrombotic microangiopathy should be assumed as a medical emergency, and treatment should be applied within the first 24 hours to guarantee safe conditions to stop and revert TMA progression.⁽¹⁸⁾ These first measures include PT, management of renal failure including evaluation of dialysis need, control of hypertension and fluid status (since hypervolemia and acute pulmonary edema are frequent complications). Red blood cells transfusions are often required to control anemia. Indication for platelet transfusion is controversial due to potential thrombotic manifestations exacerbation.^(2, 6)

The etiology may be unclear so it's important to contemplate all possible diagnosis and triggers, collect laboratory tests and prompt treatment initiation. Even though some of the results take several days (ADAMTS13 activity level, CH50, *C3*, *C4*, *FH* and *FI* levels, *FH autoantibody* level) or even months (genetic studies) to be available, it should not delay treatment initiation, as well as further adaptations according to the therapeutic response and lab results.⁽¹⁰⁾ [Fig.3](#) and [Fig.4](#) present algorithms of the management and treatment of these patients and renal transplantation approach, respectively, but there are no consensual guidelines relatively to the long-term treatment options.

PT can be supplied as plasma exchange (PE) or plasma infusion (PI). Both approaches have been crucial for mortality decrease (from 50% to 25%) in the last 3 decades.^(1, 6, 10) In PE, the patient plasma is replaced with non-native frozen plasma (FFP), with high doses of complement-regulating proteins. This exchange leads to an elimination of dysfunctional endogenous soluble complement inhibitors, *FH autoantibodies* and inflammatory/thrombogenic factors implicated in endothelial damage and platelet hyper-aggregation, explaining why PE is more effective than solely PI.^(2, 9)

Moreover PE permits a provision of larger volumes of plasma comparing with PI, avoiding fluid overload.⁽²⁾ Based on the European guidelines, PE should start with early and high volumes (1.5 to 2 plasmatic volume/day, or 60 to 75ml/kg of body weight per session) daily for 5 days, then 5 sessions/week for 2 weeks and 3 sessions/week for 2 weeks.⁽¹⁹⁾ When hemolysis became contained and platelet count stabilized PE can be gradually tapered. Patients with genetic defects usually need long-term plasma therapy (twice a week or weekly based regimens) to maintain remission.^(6, 8)

Complications of this therapy exist especially with low-weight patients (specially children) which more prone to the development of symptomatic hypervolemia.

Hypotension, allergic reactions, catheter-related complications (thrombosis and infection), and hypocalcemia are the most frequent complications with PE therapy.⁽²⁰⁾

In PI patients receive FFP and functional complement regulators (20 to 30ml/kg of body weight) but there is no replacement, so it doesn't remove the abnormal factors that modify the complement system function, reducing its efficacy.^(1, 6) Anaphylactic reactions to FFP, hypervolemia, high blood pressure, heart failure, or hyperproteinemia are the main complications.⁽⁶⁾

Therapy response (either PE or PI) varies largely according to each mutation and seems to be higher with *C3*, *thrombomodulin* mutations and *FH autoantibodies* comparing to *CFH* or *CFI* mutations.⁽¹⁰⁾ PT may induce remission in *complement factor H (CFH)* mutations, yet after long-term therapy it may become unresponsiveness. *Complement factor I (CFI)* mutations have also a partial response, with only 30 to 40% of remissions. *Complement factor B* and *C3* mutations achieve rates of 30 and 50% of response respectively, but patients carrying these mutations will need more regular plasma exchanges to eliminate the hyper-functional mutant proteins.⁽²⁾

CFH autoantibodies are cleared in PE so this therapy can be used to improve the results in these patients since high antibody titers are associated with a higher risk of relapse. The rational for PE combination with immunosuppressive agents, in this group of patients, is justified by the reduction of antibody production.⁽⁶⁾

MCP is a transmembrane glycoprotein, surface-bound complement regulatory protein that acts as a cofactor for the *CFI*-mediated cleavage of *C3b* and *C4b* that are deposited on host cells. Patients with *MCP* mutations carry the best prognosis albeit plasma treatment did not impact the outcome significantly: remission was achieved in around 90% of both plasma-treated and plasma-untreated acute episodes. Renal transplantation

outcome is more favorable in patients with *MCP* mutations, than in patients with *CFH* and *CFI* mutations due to disease recurrence in the last ones.^(2, 10)

The clinical response after 5-7 days of daily PT treatment should aid further decisions since the complete hematological response and renal recovery rates are usually below 50% and differ according to the type of mutations or the levels of antibody titers.^(5, 9)

The explanation for that low recovery rate relies on the facts that PT doesn't inhibit the underlying complement-mediated pathogenic mechanism or prevent progression of tissue damage.⁽²¹⁾

Patients with cancer are an exception since PT will not improve the outcome in these cases, highlighting the importance of the neoplastic disease control.⁽¹⁸⁾

The American Society for Apheresis (ASFA) present guidelines with plasma therapy indications, which are graded from I to IV, being category I a 1st line indication to PT based on the evidence and category IV when the evidence shows ineffective results. Since aHUS due to *CFH autoantibodies* fits in the category I, early PT is vital on the syndrome management.⁽⁵⁾ In cases of incomplete response, maintenance of MAHA and absence of renal recovery eculizumab may be considered.^(5, 8) However, the most effective approach is currently not consensual.⁽⁶⁾

Eculizumab is a recombinant humanized monoclonal antibody that by a high affinity binding to *C5* blocks the cleavage into its effectors *C5a* and *C5b* and stops the membrane attack complex formation.⁽²²⁾ Regardless the type of mutation or complement anomaly eculizumab can be effective, given that *C5* is a terminal complement component.⁽¹⁾ However, recent studies showed a residual activity of the terminal pathway through alternative pathway activation and elevated *C5a* levels in patients

treated with eculizumab, since it fails to block *C5a* resulting from direct enzyme cleavage by trypsin and thrombin .^(23, 24)

Extensive trials, prospective studies and case reports demonstrated the effectiveness of eculizumab, reporting that about 85% of the patients became disease-free, even in those plasma-resistant.^(6, 8, 25) Results in long-term also showed a stabilization of the platelet count with a decrement no more than 25% from the baseline, allowing PT discontinuation. Moreover, improvement of renal function with dialysis suspension in some cases was also observed, as well as a higher quality of life, safety and tolerability.^(1, 26) Eculizumab also demonstrate satisfactory results relatively to complement-related, inflammation and endothelial activation biomarkers, even though some of them did not completely normalize, emphasizing the constant risk of chronic activation of complement system in aHUS patients.⁽⁶⁾ However, for aHUS patients carrying *DGK ϵ* mutation eculizumab doesn't seem to be effective.⁽¹¹⁾ Familial history of HUS, relapsed aHUS, high-risk genotypes, renal transplantation, resistance or dependency on plasma therapy are factors that favor the use of eculizumab.⁽¹⁰⁾ Furthermore treatment with eculizumab should started be as early as possible, soon after STEC-HUS and TTP diagnosis exclusion.⁽²⁷⁾ Note that some trials demonstrate a correlation inversely proportional between the improvement of the estimated glomerular filtration rates and the time between the clinical manifestations and the beginning of the drug.⁽⁶⁾ The recommended dose in adults is 900mg per week intravenous infusion, for 4 weeks (induction phase), followed by a 5th dose of 1200mg after 7 days, and afterwards 1200mg every 14 days (maintenance phase).⁽²⁸⁾ Patients who are also receiving PE should do an extra dose of 600mg of eculizumab after each plasma session.⁽¹⁰⁾ The length of the treatment it's not clearly established but cumulative experience may facilitate consensus in treatment strategies.⁽²⁷⁾ The assessment of response to

eculizumab is based on monitoring disease activity markers like platelet counts, lactate dehydrogenase, haptoglobin, serum creatinine and complement activity markers.⁽²⁹⁾ When complement inhibition is successful, it's observed considerable reductions of the hemolytic complement function: the classical (CH50), the alternative pathway (APH50), and the *C3d* and *sC5b-9* levels, while *C5a* levels diminished only 6 months after eculizumab. Drug accumulation can be demonstrated by serum concentration of eculizumab (level over to 1082 µg/ml). Specific enzyme-linked immunosorbent assay (ELISA) is also a way to determine eculizumab concentration.⁽³⁰⁾ Long-term treatment is recommended in renal transplantation, in high-risk genotypes, and in severe extra-renal findings. Several authors suggest life-long treatment with eculizumab, due to its the favorable side-effect profile.^(8, 31)

Some studies reported eculizumab deposition in glomeruli with anti-eculizumab antibody production but the long-term significance is unclear.⁽³²⁾

The decision for eculizumab withdrawal must take in account the risk of recurrence, the type of mutation, the patient age, the renal function and extra-renal manifestations. The patient must be closely monitored and in the presence of relapse, re-administration should be prompt initiated.⁽³³⁾ Regular self-monitoring of hemoglobinuria through urine dipsticks has been recently introduced as a form to quickly detect aHUS relapses after discontinuation of eculizumab.⁽³⁴⁾

Treatment with eculizumab has an annual cost of 700,000\$ approximately and laboratory assays to optimize the treatment are lacking, so Kerboua. K, et al suggested a novel disease activity marker, C3:CH50 ratio, to use during the induction and maintenance phase.^(35, 36)

Because of the augmented risk of infection by encapsulated organisms of patients under eculizumab treatment, tetravalent vaccine against *Neisseria meningitidis* (A, C, Y, W135) as well as vaccination against serogroup B two weeks before eculizumab treatment is mandatory. After that, patients should receive prophylactic antibiotics (penicillin V potassium at 250 mg every 12 hours or ciprofloxacin 500 mg daily during 4 weeks).⁽³⁷⁾ At least two months after the first vaccine a booster dose should be performed.⁽¹⁰⁾ When the postponement of eculizumab treatment until vaccine response isn't possible associated treatment with antibiotics against *N. meningitides* may be initiated and antibiotic prophylaxis may be instituted according the hospital protocol.⁽⁶⁾ To immunocompromised patients it should be given continued antibiotic prophylaxis as long as eculizumab treatment last.⁽³⁷⁾ Vaccination against *Haemophilus influenzae* type b is recommended annually to all patients and vaccination against *Pneumococcus pneumoniae* should also be considered in children case.^(5, 10) During active infections, eculizumab should be suspended if possible. Counseling and screening for gonococcal infection should be performed after and during treatment in sexually active patients, mainly in high risk patients, and in their partners.⁽³⁷⁾

The most common adverse effects of eculizumab include cough, headache, nausea, vomiting, diarrhea, hypertension, upper respiratory tract infections, urinary tract infections and abdominal pain.⁽¹⁾ Hepatotoxicity, a pediatric side-effect, was also described in an adult patient.⁽³⁸⁾

Despite the best results with eculizumab, this treatment costs 4800-6500€ per 300mg, being four times more expensive than PE (850-1800€ per session) and making it inaccessible to many low-resource settings.⁽³⁹⁾

The outcome of renal transplantation in aHUS patients is variable, having a straight correlation with the genetic abnormality involved as shown on Tab.1.⁽⁴⁰⁾ For instance

CFH, *CFI*, *C3* and *CFB* mutation carriers have abnormal complement regulators, most of them synthesized in the liver, with high recurrence rate, reaching up to 80% of the cases.⁽⁴¹⁾ On the other hand, *MCP* mutations carries more favorable outcomes, with free-recurrence rates of 80%, being suggested that the individuals who recur have an additional genetic predisposition.⁽⁸⁾ These good results can be explained because *MCP* is a transmembrane protein highly expressed on kidney, so the defect is corrected when kidney transplant is performed.⁽⁴²⁾

Post-transplant aHUS recurrence occurs in 60 to 90% in the first year after transplant and graft-survival is only 51%.⁽⁴³⁾ It may be triggered by brain-death related injury, ischemia-reperfusion injury, infections (cytomegalovirus, influenza virus, parvovirus B19, BK virus, upper respiratory tract infections and infectious gastroenteritis), the use of immunosuppressive drugs (calcineurin inhibitors, mTOR inhibitors) or rejection.⁽⁴⁴⁾

There are limited data on the efficacy of prophylactic plasma therapy including plasmapheresis to prevent post-transplant aHUS recurrence. Nevertheless, the progress in genetic studies and the introduction of eculizumab allowed renal transplantation in highly risk patients.^(1, 8) Albeit the promising results published in some living donor transplantation cases with prophylactic eculizumab regimens, living donation isn't consensual. The risk for the presence of mutations or polymorphisms still not detectable in the potential living donors is a concern and may justify the cases of aHUS in healthy donors, probably triggered after the donation procedure.⁽⁴³⁾

Beyond PE and eculizumab, there are other therapeutic options in order to avoid recurrence after renal transplantation such as liver-kidney combined transplantation and the use of antibody-suppressing or depleting agents (corticosteroids, azathioprine, mycophenolate mofetil, cyclophosphamide or rituximab). These agents have shown

important results relatively to *CFH autoantibodies* control, especially with rituximab. The absence of aHUS recurrence, normalized graft function and negative titers in previous studies using rituximab, turned it into a beneficial prophylactic choice to patients with persistently elevated *CFH autoantibodies* titers, even when renal transplant is not performed.^(10, 42)

Recombinant-human C1 inhibitor (C1-INH) is being tested as therapy for antibody-mediated rejection and to prevent ischemia-reperfusion injury.⁽⁴¹⁾

Since complement plasma factors like *FH*, *FI*, *FB* and *C3* are mainly synthesized in the liver, it seemed logical that combined renal and liver transplantation could correct the genetic deficiency and so normalize the activity of fluid-phase complement regulatory proteins.^(6, 8) However, the results were not so promising, leading to ischemia-reperfusion injuries, post-transplant recurrence and fatal encephalopathy, with 10-20% of short-term mortality.^(2, 6, 8) The high morbidity and mortality associated to this procedure, and the organ shortage, make this option a rare indication.⁽⁶⁾ When contemplated, screening for mutations, prophylactic eculizumab or PE, in addition to aspirin and heparin, must be performed.^(2, 6, 8)

Calcineurin (CNI)-free regimens has been attempted in a few reports, based in Belatacept regimens, with favorable short-term outcomes.⁽⁴²⁾ The rationale for those protocols aim to reduce the introduction of potential drug triggers.

The treatment of post-transplant recurrence consists in early initiation of PE or eculizumab.⁽⁴²⁾ However, new treatments and approaches are arising in the aHUS field.

SKY59, a novel humanized antibody has a long-lasting neutralization of C5 activity comparing to eculizumab.⁽⁴⁵⁾ Studies with recombinant factor H also demonstrated optimistic results in the treatment of patients with CFH deficiency.^(1, 2) Inhibitors of

angiotensin-converting enzyme, statins (HMG-CoA reductase), ascorbic acid and allopurinol (inhibitor of xanthine oxidase) are also potential adjuvant agents for aHUS treatment since they reduce the oxidative inactivation of nitric oxide, increasing nitric oxide availability and promoting the endothelium-dependent vasodilation.⁽¹⁾ Other new options are the oral CCX168 (C5aR antagonist), currently in a phase 2 trial, and a small-molecule factor D inhibitor that successfully inhibited alternative complement pathway by binding human Factor D and inhibiting its proteolytic activity against purified Factor B in complex with C3b.^(5, 46, 47)

EVOLUTION AND PROGNOSIS

aHUS is prone to a poor prognosis, due to the underlying genetic defect, whose recognition is essential for risk stratification after renal transplantation as shown on Tab-1.⁽⁸⁾ ESRD, extra-renal symptoms, number of recurrent episodes after renal transplantation and the delay time until prompt initiation therapy, also change the prognosis.⁽¹⁰⁾ Despite that, aHUS prognosis is reserved with high rates of mortality (25% in the acute phase), end-stage renal disease (50% within a year) and recurrence.⁽¹⁾

CONCLUSION

The perspectives of treatment aHUS changed drastically in the past few years, after eculizumab introduction in the therapeutic armamentarium, providing a useful tool in aHUS PE resistant or dependent, as well in renal transplantation. Nevertheless, the cost remains a major obstacle for its generalized implementation, leaving PE as the only therapeutic option available in many centers. Moreover, consensual guidelines with precise eculizumab indications and management strategies are needed in order to provide adequate follow-up of this diverse group of patients.

Further information about the genetic and the pathophysiology of aHUS is crucial for the development of therapeutic strategies and, as a consequence, new therapeutics molecules acting on specific targets in complement system may provide promising results in the near future.

ACKNOWLEDGEMENTS

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

REFERENCES:

1. Nayer A, Asif A. Atypical Hemolytic-Uremic Syndrome: A Clinical Review. *Am J Ther*. 2016;23(1):e151-8.
2. Noris M, Remuzzi G. Atypical hemolytic-uremic syndrome. *N Engl J Med*. 2009;361(17):1676-87.
3. George JN, Nester CM. Syndromes of thrombotic microangiopathy. *N Engl J Med*. 2014;371(19):1847-8.
4. Noris M, Mescia F, Remuzzi G. STEC-HUS, atypical HUS and TTP are all diseases of complement activation. *Nat Rev Nephrol*. 2012;8(11):622-33.
5. Zhang K, Lu Y, Harley KT, Tran MH. Atypical Hemolytic Uremic Syndrome: A Brief Review. *Hematol Rep*. 2017;9(2):7053.
6. Campistol JM, Arias M, Ariceta G, Blasco M, Espinosa L, Espinosa M, et al. An update for atypical haemolytic uraemic syndrome: diagnosis and treatment. A consensus document. *Nefrologia*. 2015;35(5):421-47.
7. Goodship TH, Cook HT, Fakhouri F, Fervenza FC, Fremeaux-Bacchi V, Kavanagh D, et al. Atypical hemolytic uremic syndrome and C3 glomerulopathy:

conclusions from a "Kidney Disease: Improving Global Outcomes" (KDIGO) Controversies Conference. *Kidney Int.* 2017;91(3):539-51.

8. Kavanagh D, Goodship TH, Richards A. Atypical hemolytic uremic syndrome. *Semin Nephrol.* 2013;33(6):508-30.

9. Brocklebank V, Wood KM, Kavanagh D. Thrombotic Microangiopathy and the Kidney. *Clin J Am Soc Nephrol.* 2017.

10. Afshar-Kharghan V. Atypical hemolytic uremic syndrome. *Hematology Am Soc Hematol Educ Program.* 2016;2016(1):217-25.

11. Jokiranta TS. HUS and atypical HUS. *Blood.* 2017;129(21):2847-56.

12. Loirat C, Fremeaux-Bacchi V. Atypical hemolytic uremic syndrome. *Orphanet J Rare Dis.* 2011;6:60.

13. Nester CM, Smith RJ. Factors influencing treatment of atypical hemolytic uremic syndrome. *Clin J Am Soc Nephrol.* 2014;9(9):1516-8.

14. Jin A, Boroujerdi-Rad L, Shah G, Chen JL. Thrombotic microangiopathy and human immunodeficiency virus in the era of eculizumab. *Clin Kidney J.* 2016;9(4):576-9.

15. Shen YM. Clinical evaluation of thrombotic microangiopathy: identification of patients with suspected atypical hemolytic uremic syndrome. *Thromb J.* 2016;14(Suppl 1):19.

16. Huerta A, Arjona E, Portoles J, Lopez-Sanchez P, Rabasco C, Espinosa M, et al. A retrospective study of pregnancy-associated atypical hemolytic uremic syndrome. *Kidney Int.* 2018;93(2):450-9.

17. Karpman D, Loos S, Tati R, Arvidsson I. Haemolytic uraemic syndrome. *J Intern Med.* 2017;281(2):123-48.

18. Azoulay E, Knoebl P, Garnacho-Montero J, Rusinova K, Galstian G, Eggimann P, et al. Expert Statements on the Standard of Care in Critically Ill Adult Patients With Atypical Hemolytic Uremic Syndrome. *Chest*. 2017;152(2):424-34.
19. Bajracharya P, Jain A, Baracco R, Mattoo TK, Kapur G. Atypical hemolytic uremic syndrome: a clinical conundrum. *Pediatr Nephrol*. 2016;31(10):1615-24.
20. Palma LM, Langman CB. Critical appraisal of eculizumab for atypical hemolytic uremic syndrome. *J Blood Med*. 2016;7:39-72.
21. Legendre CM, Licht C, Muus P, Greenbaum LA, Babu S, Bedrosian C, et al. Terminal complement inhibitor eculizumab in atypical hemolytic-uremic syndrome. *N Engl J Med*. 2013;368(23):2169-81.
22. Cavero T, Rabasco C, Lopez A, Roman E, Avila A, Sevillano A, et al. Eculizumab in secondary atypical haemolytic uraemic syndrome. *Nephrol Dial Transplant*. 2017;32(3):466-74.
23. Puissant-Lubrano B, Puissochet S, Congy-Jolivet N, Chauveau D, Decramer S, Garnier A, et al. Alternative complement pathway hemolytic assays reveal incomplete complement blockade in patients treated with eculizumab. *Clin Immunol*. 2017;183:1-7.
24. Riedemann NC, Habel M, Ziereisen J, Hermann M, Schneider C, Wehling C, et al. Controlling the anaphylatoxin C5a in diseases requires a specifically targeted inhibition. *Clin Immunol*. 2017;180:25-32.
25. Licht C, Greenbaum LA, Muus P, Babu S, Bedrosian CL, Cohen DJ, et al. Efficacy and safety of eculizumab in atypical hemolytic uremic syndrome from 2-year extensions of phase 2 studies. *Kidney Int*. 2015;87(5):1061-73.
26. Yamada Y, Abe R, Okano Y, Miyakawa Y. Long-term Eculizumab Treatment Contributes to Recovery from End-stage Renal Disease Caused by Atypical Hemolytic Uremic Syndrome. *Intern Med*. 2017;56(9):1085-8.

27. Ebrahem R, Kadhem S, Truong Q. Treatment of Atypical Hemolytic-Uremic Syndrome in the Era of Eculizumab. *Cureus*. 2017;9(3):e1111.
28. Volokhina E, Wijnsma K, van der Molen R, Roeleveld N, van der Velden T, Goertz J, et al. Eculizumab Dosing Regimen in Atypical HUS: Possibilities for Individualized Treatment. *Clin Pharmacol Ther*. 2017;102(4):671-8.
29. Ardissino G, Tel F, Sgarbanti M, Cresseri D, Giussani A, Griffini S, et al. Complement functional tests for monitoring eculizumab treatment in patients with atypical hemolytic uremic syndrome: an update. *Pediatr Nephrol*. 2018;33(3):457-61.
30. Wehling C, Amon O, Bommer M, Hoppe B, Kentouche K, Schalk G, et al. Monitoring of complement activation biomarkers and eculizumab in complement-mediated renal disorders. *Clin Exp Immunol*. 2017;187(2):304-15.
31. Rodriguez E, Barrios C, Soler MJ. Should eculizumab be discontinued in patients with atypical hemolytic uremic syndrome? *Clin Kidney J*. 2017;10(3):320-2.
32. Carter S, Hewitt I, Kausman J. Long-term remission with eculizumab in atypical haemolytic uraemic syndrome. *Nephrology (Carlton)*. 2017;22 Suppl 1:7-10.
33. Sahutoglu T, Basturk T, Sakaci T, Koc Y, Ahbap E, Sevinc M, et al. Can eculizumab be discontinued in aHUS?: Case report and review of the literature. *Medicine (Baltimore)*. 2016;95(31):e4330.
34. Fakhouri F, Fila M, Provot F, Delmas Y, Barbet C, Chatelet V, et al. Pathogenic Variants in Complement Genes and Risk of Atypical Hemolytic Uremic Syndrome Relapse after Eculizumab Discontinuation. *Clin J Am Soc Nephrol*. 2017;12(1):50-9.
35. Kerboua KE, Haiba F, Batouche D. C3:CH50 ratio as a proposed composite marker for eculizumab monitoring in atypical hemolytic uremic syndrome: Preliminary results. *J Immunoassay Immunochem*. 2017;38(2):178-89.

36. Noris M, Ruggenenti P, Remuzzi G. Kidney Transplantation in Patients With Atypical Hemolytic Uremic Syndrome: A Therapeutic Dilemma (or Not)? *Am J Kidney Dis.* 2017;70(6):754-7.
37. Benamu E, Montoya JG. Infections associated with the use of eculizumab: recommendations for prevention and prophylaxis. *Curr Opin Infect Dis.* 2016;29(4):319-29.
38. Oruc A, Ayar Y, Vuruskan BA, Yildiz A, Aktas N, Yavuz M, et al. Hepatotoxicity associated with eculizumab in a patient with atypical hemolytic uremic syndrome. *Nefrologia.* 2017.
39. Keenswijk W, Walle JV. Atypical Hemolytic Uremic Syndrome in Low Resource Settings: Which Options Do We Have? *Ther Apher Dial.* 2017.
40. Milan Manani S, Virzi GM, Giuliani A, Clementi A, Brocca A, Dissegna D, et al. Hemolytic Uremic Syndrome and Kidney Transplantation: A Case Series and Review of the Literature. *Nephron.* 2017;136(3):245-53.
41. Grenda R, Durlik M. Eculizumab in Renal Transplantation: A 2017 Update. *Ann Transplant.* 2017;22:550-4.
42. Noris M, Remuzzi G. Managing and preventing atypical hemolytic uremic syndrome recurrence after kidney transplantation. *Curr Opin Nephrol Hypertens.* 2013;22(6):704-12.
43. de Andrade LGM, Contti MM, Nga HS, Bravin AM, Takase HM, Viero RM, et al. Long-term outcomes of the Atypical Hemolytic Uremic Syndrome after kidney transplantation treated with eculizumab as first choice. *PLoS One.* 2017;12(11):e0188155.
44. Okumi M, Tanabe K. Prevention and treatment of atypical haemolytic uremic syndrome after kidney transplantation. *Nephrology (Carlton).* 2016;21 Suppl 1:9-13.

45. Fukuzawa T, Sampei Z, Haraya K, Ruike Y, Shida-Kawazoe M, Shimizu Y, et al. Long lasting neutralization of C5 by SKY59, a novel recycling antibody, is a potential therapy for complement-mediated diseases. *Sci Rep*. 2017;7(1):1080.
46. Lorthiois E, Anderson K, Vulpetti A, Rogel O, Cumin F, Ostermann N, et al. Discovery of Highly Potent and Selective Small-Molecule Reversible Factor D Inhibitors Demonstrating Alternative Complement Pathway Inhibition in Vivo. *J Med Chem*. 2017;60(13):5717-35.
47. Yuan X, Gavrilaki E, Thanassi JA, Yang G, Baines AC, Podos SD, et al. Small-molecule factor D inhibitors selectively block the alternative pathway of complement in paroxysmal nocturnal hemoglobinuria and atypical hemolytic uremic syndrome. *Haematologica*. 2017;102(3):466-75.

TABLES AND FIGURES:


Table 1. Atypical hemolytic uremic syndrome: genetic modifications, clinical characteristics and prognosis.

		Factor H	Factor I	MCP – membrane cofactor protein	Complement C3	Factor B	Thrombomodulin	FH autoantibodies
Gene mutation		CFH	CFI	CD46	C3	CFB	THBD	CFHR1-3 deletion
Location of the mutation		Chr 1q32	Chr 4q25	Chr 1q32	Chr 19p13.3	Chr 6p21.3	Chr 20p11.2	Chr1
Frequency (%)		24-28	4-8	5-9	2-8	0-4	0-5	3-10
Minimal age onset	Children	Birth	Birth	> 1 year	7 months	1 month	6 months	Mostly 7-11 years
	Adult	Any age	Any age	Any age	Any age	Any age	rare	
Clinical presentation				Typically present in childhood, milder acute episode	Typically present in childhood	Higher variability	90% present in childhood	Significant gastrointestinal symptoms and/or diarrhea thus, resembling aHUS
Risk of relapse (%)		20-40	20-40	80	20-40	70	30	60* 10†
Relapse after RT (%) ¥		80-90	70-80	15-20	40-50	88	-	20
Death or ESRD in the first episode or within > 1 year (%)		50-70	50	0-6	60	50	50	30-40
Death or ESRD 3-10 years after onset (%)		70-80	60-70	<20	60-70	70	50-60	30-70
Risk of discontinue eculizumab		high	low	low	high			Low for homozygous deletion

RT: renal transplantation; ESRD: End-stage renal disease; ¥Without prophylactic therapy

*Without immunosuppression; †With immunosuppression


Figure 1. The alternative pathway of the complement system and the action of eculizumab.



The alternative pathway (AP) of the complement system is continuously activated low-activated in plasma, finishing with the formation of the membrane-complex attack (MAC) and with the release of powerful anaphylotoxins (C5a). In the beginning of the process, C3(H₂O) binds factor B (CFB), which is cleaved by factor D (CFD), and form the alternative pathway initiation C3 convertase. This is responsible to cleave C3 into C3a and C3b. C3b and hydroxyl groups on cell-surface carbohydrates and proteins of target cells establish a covalent binding forming the amplification loop C3 convertase. This C3b also binds CFB and form another amplification loop C3 convertase (on the fluid phase). Cleavage of C3 molecules by C3 convertase enzymes ends on another feedback amplification loop. At least, C3b binds to C3bBb creating the C5 convertase enzyme, responsible to cleave C5 into C5a and C5b which will be part of the MAC. This mechanism is extremely regulated by complement factor H (CFH), complement factor I (CFI), complement receptor 1 (CR1), decay accelerating factor (DAF) and membrane cofactor protein (MCP) in a way to prevent host cell injury and control


the deposition of complement in pathogens cell surface. CR1 and CFH both perform as cofactors for CFI, leading to a cleavage of C3b and a decrease of the C3 convertase. MCP binds to C3b preventing its connection to other factors and DAF is responsible to dissociate C3 and C5 convertases. Eculizumab is a monoclonal antibody that that by a high affinity binding to C5 blocks the cleavage into its effectors C5a and C5b and stops the membrane attack complex formation.

Figure 2. Cycle and parallel processes and the action on the cells of the complement system activation in aHUS.



The dysregulated activation of the complement system happens where there are abnormal regulator proteins (CFH, CFI or MCP) or overactive mutated complement factors (CFB, C3), leading to the release of anaphylotoxins (C5a) and to the membrane attack complex (MAC) formation. In red cells, this ends with release of the hemoglobin and decrease of the nitric oxide (NO) in plasma. Platelet activation and aggregation and augmented tissue factor (TF) activity on the endothelium are also a consequence of this complement processes. These mechanisms lead to a procoagulative state, coagulation and tissue impairment. The feedback loops observed could explain the cyclical and parallel characteristics of this phenomena. Thrombomodulin, diacylglycerol kinase ϵ (DGK ϵ) and plasmin are also responsible to control these mechanism, and abnormalities in these proteins can also be an explanation to the pathogenic processes in aHUS.

Figure 3. Diagnosis and treatment algorithm for the management of a patient with thrombotic microangiopathy (TMA)




Supportive treatment is mandatory and includes, besides fluid management, red blood cells transfusion, renal replacement therapy and intensive care unit admission if necessary. Platelet transfusion should be avoided since it could aggravate the TMA. TMA approach should start to exclude TTP and STEC-HUS by ADAMST13 activity assays and Stx detection, respectively. Because of the high mortality rates of TTP if untreated, early plasma therapy (PT) should be

instituted in adults but since TTP is rare in children, first line treatment could be eculizumab if aHUS is suspected. A complete clinical history should also be collected to establish potential triggers/causes. Evidence suggesting a complement etiology could include familiar history or previous aHUS episodes and in those cases we could perform a complement study, stop PT and start eculizumab if possible. If eculizumab isn't available, continuation of PT therapy or renal transplant could be considered. When the patient has elevated FH antibody titers, a combined therapy with rituximab or other immunosuppressive agent could be contemplated.

ADAMTS13: a disintegrin and metalloproteinase with a thrombospondin type 1 motif, member 13; aHUS: atypical hemolytic uremic syndrome; DGKE: gene encoding diacylglycerol kinase; CFH: complement factor H; CFI: complement factor I; CFB: complement Factor B; HUS: hemolytic uremic syndrome; LDH: lactate dehydrogenase; MCP: membrane cofactor protein; PE: plasma exchange; PT: Plasma therapy; STEC: shiga toxin-producing *Escherichia coli*; Stx: shiga toxin; THBD: Thrombomodulin; TMA: thrombotic microangiopathy; TTP: thrombotic thrombocytopenic purpura.

Figure 4. Atypical hemolytic uremic syndrome management algorithm for patients eligible for renal transplantation.



ESRD: end-stage renal disease; MCP: membrane cofactor protein; DGKE: diacylglycerol kinase ϵ ; FH: Factor H; CFH: complement factor H; CFI: complement factor I; CFB: complement factor B; aHUS: atypical hemolytic uremic syndrome; RT: renal transplantation; PT: plasma therapy (in these plasma exchange should be performed). * Living donor transplantation is controversial and should only be considered when eculizumab is available.



SEMINARS IN NEPHROLOGY

AUTHOR INFORMATION PACK

TABLE OF CONTENTS

●	Description	p.1
●	Impact Factor	p.1
●	Abstracting and Indexing	p.1
●	Editorial Board	p.1
●	Guide for Authors	p.2



ISSN: 0270-9295

DESCRIPTION

Seminars in Nephrology is a timely source for the publication of new concepts and research findings relevant to the clinical practice of **nephrology**. Each issue is an organized compendium of practical information that serves as a lasting reference for nephrologists, internists and physicians in training.

Benefits to authors

Elsevier provides many author benefits, such as free PDFs, a liberal copyright policy, special discounts on Elsevier publications and much more. Please click here for more information on our [author services](#).

If you require any further information or help, please visit our [Support Center](#)

IMPACT FACTOR

2016: 3.598 © Clarivate Analytics Journal Citation Reports 2017

ABSTRACTING AND INDEXING

MEDLINE®
Current Contents/Clinical Medicine
Science Citation Index
EMBASE

EDITORIAL BOARD

Editor

Joseph V. Bonventre, MD, PhD, Samuel A. Levine Professor of Medicine; Chief, Renal Division; and Chief, Division of Engineering in Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

Editorial Advisory Board

Glenn Chertow, MD, San Francisco
Benjamin Humphreys, MD, PhD, Boston
Adeera Levin, MD, Vancouver
John Sedor, MD, Cleveland
Stuart Shankland, MD, Seattle
Myles Wolf, MD, Miami

GUIDE FOR AUTHORS

INTRODUCTION

Seminars in Nephrology is a topical journal that provides scholarly review articles focusing on subjects of current importance in clinical nephrology, kidney research and related fields. Each issue examines a single topic and is meant to keep the practicing nephrologist up to date on the clinical relevance of new investigations and the scientific underpinnings of nephrology. The journal covers all aspects of the clinical practice of nephrology.

Submission Information

Each issue of *Seminars in Nephrology* is devoted to a specific topic. The Guest Editor of each issue invites contributors who are experts in the field to submit manuscripts. Invited authors will find manuscript preparation guidelines and other publishing information in this document. Suggestions for future topics are welcome and may be sent to the Editor

Deadline

It is of utmost importance to submit your manuscript, complete with all figures and tables (and written permission to reproduce them when required), to the Guest Editor by the specified deadline. This deadline provides time for the Guest Editor and Dr. Bonventre to review your manuscript and forward it to the publisher in time for prompt publication. It will also allow time for any revisions to your manuscript should they be necessary. Please contact the Guest Editor if you are unsure about the deadline.

BEFORE YOU BEGIN

Ethics in publishing

Please see our information pages on [Ethics in publishing](#) and [Ethical guidelines for journal publication](#).

Conflict of Interest

All authors must disclose on the title page of their article any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding.

Submission declaration

Submission of an article implies that the work described has not been published previously (except in the form of an abstract, a published lecture or academic thesis, see '[Multiple, redundant or concurrent publication](#)' for more information), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder.

Preprints

Please note that [preprints](#) can be shared anywhere at any time, in line with Elsevier's [sharing policy](#). Sharing your preprints e.g. on a preprint server will not count as prior publication (see '[Multiple, redundant or concurrent publication](#)' for more information).

Copyright

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (see [more information](#) on this). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. [Permission](#) of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations. If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has [preprinted forms](#) for use by authors in these cases.

For open access articles: Upon acceptance of an article, authors will be asked to complete an 'Exclusive License Agreement' ([more information](#)). Permitted third party reuse of open access articles is determined by the author's choice of [user license](#).

Author rights

As an author you (or your employer or institution) have certain rights to reuse your work. [More information](#).

Elsevier supports responsible sharing

Find out how you can [share your research](#) published in Elsevier journals.

Role of the funding source

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated.

Funding body agreements and policies

Elsevier has established a number of agreements with funding bodies which allow authors to comply with their funder's open access policies. Some funding bodies will reimburse the author for the Open Access Publication Fee. Details of [existing agreements](#) are available online.

After acceptance, open access papers will be published under a noncommercial license. For authors requiring a commercial CC BY license, you can apply after your manuscript is accepted for publication.

Open access

This journal offers authors a choice in publishing their research:

Subscription

- Articles are made available to subscribers as well as developing countries and patient groups through our [universal access programs](#).
- No open access publication fee payable by authors.

Open access

- Articles are freely available to both subscribers and the wider public with permitted reuse.
- An open access publication fee is payable by authors or on their behalf, e.g. by their research funder or institution.

Regardless of how you choose to publish your article, the journal will apply the same peer review criteria and acceptance standards.

For open access articles, permitted third party (re)use is defined by the following [Creative Commons user licenses](#):

Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

For non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.

The open access publication fee for this journal is **USD 2500**, excluding taxes. Learn more about Elsevier's pricing policy: <http://www.elsevier.com/openaccesspricing>.

Green open access

Authors can share their research in a variety of different ways and Elsevier has a number of green open access options available. We recommend authors see our [green open access page](#) for further information. Authors can also self-archive their manuscripts immediately and enable public access from their institution's repository after an embargo period. This is the version that has been accepted for publication and which typically includes author-incorporated changes suggested during submission, peer review and in editor-author communications. Embargo period: For subscription articles, an appropriate amount of time is needed for journals to deliver value to subscribing customers before an article becomes freely available to the public. This is the embargo period and it begins from the date the article is formally published online in its final and fully citable form. [Find out more](#).

This journal has an embargo period of 12 months.

Elsevier Researcher Academy

[Researcher Academy](#) is a free e-learning platform designed to support early and mid-career researchers throughout their research journey. The "Learn" environment at Researcher Academy offers several interactive modules, webinars, downloadable guides and resources to guide you through the process of writing for research and going through peer review. Feel free to use these free resources to improve your submission and navigate the publication process with ease.

Language (usage and editing services)

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the [English Language Editing service](#) available from Elsevier's WebShop.

Submission

Please send your manuscript and figures, and letters of permission for previously published figures and tables to the Guest Editor who has invited you to contribute. Manuscripts and figures should be sent electronically. Figures submitted electronically should be in a separate file.

PREPARATION

PREPARATION

All components of the electronic manuscript must appear within a single electronic file: references, figure legends, and tables must appear at the end of the manuscript.

Your manuscript should consist of the following elements, each starting on a separate page: • Title page • Abstract and key words • Body of manuscript • References • Figure legends • Figures (with permissions for reuse, if required) • Tables (with permissions for reuse, if required) • Supplemental materials for online-only publication, if desired

All parts of the manuscript (including references and legends) should be prepared double-spaced, with a full line of space after every line and with 1.5-inch margins on both sides and at the top and bottom of every page. Your abstract should not be more than 200 words. Key words should be a list of three to five important words or phrases for indexing.

All DRUG NAMES cited in the manuscript should be generic, followed by brand name, manufacturer, city, and state (or country, if not U.S.) in parentheses.

Please explain all abbreviations in parentheses the first time they are used in the text.

ESTIMATING MANUSCRIPT LENGTH

Please adhere to the estimated number of printed pages the Guest Editor has specified for your article. We are counting on your cooperation in remaining within this allocation and within our budget of 80 printed pages per issue for Seminars in Nephrology. Please contact the Guest Editor if you are unsure of the appropriate length for your article. As a rule, your manuscript should have at least 20 double-spaced pages of text and not more than 25 double-spaced pages, excluding references.

Figures and tables must total at least 4 for the submission. Your total number of references should not exceed 100.

The following approximations are offered to help you calculate the appropriate length of your manuscript: 3 8 ½ x 11-inch pages, double-spaced, with 1.5-inch margins, in standard 10- or 12-point word processor type (approx. 250 words per page) = 1 printed page 35 references (all lines double-spaced) = 1 printed page 4 tables or figures with legends = 1 printed page

Example: A manuscript contains 21 pages of text with 35 references and 4 tables: 21 pages of text = 7 printed pages 4 figures or tables = 1 printed page 35 references = 1 printed page Total = 9 printed pages

Subdivision - unnumbered sections

Divide your article into clearly defined sections. Each subsection is given a brief heading. Each heading should appear on its own separate line. Subsections should be used as much as possible when cross-referencing text: refer to the subsection by heading as opposed to simply 'the text'.

Essential Title Page Information

On the TITLE PAGE, please include all of the following information: 1. The names, degrees, and professional affiliations (position, department, institution, place) of all authors. 2. The name of the institution where the work reported was done ("From..."). 3. Acknowledgement of grant support when appropriate ("Supported in part by..."). 4. The corresponding author's telephone and fax numbers, email address, and complete mailing address (including U.S. zip code or other countries' postal code), for reprint requests and other correspondence.

Abstract

The abstract should state briefly the status of the field being reviewed, the importance to clinical medicine and exciting and novel new directions integrating science and clinical nephrology. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, references should be avoided, but if essential, then cite the author(s) and year(s). Also, nonstandard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

Keywords

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

Abbreviations

Define abbreviations that are not standard in this field in a footnote to be placed on the first page of the article. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

Acknowledgements

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here funding sources as well as those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

Formatting of funding sources

List funding sources in this standard way to facilitate compliance to funder's requirements:

Funding: This work was supported by the National Institutes of Health [grant numbers xxxx, yyyy]; the Bill & Melinda Gates Foundation, Seattle, WA [grant number zzzz]; and the United States Institutes of Peace [grant number aaaa].

It is not necessary to include detailed descriptions on the program or type of grants and awards. When funding is from a block grant or other resources available to a university, college, or other research institution, submit the name of the institute or organization that provided the funding.

If no funding has been provided for the research, please include the following sentence:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Units

Follow internationally accepted rules and conventions: use the international system of units (SI). If other units are mentioned, please give their equivalent in SI.

Math formulae

Please submit math equations as editable text and not as images. Present simple formulae in line with normal text where possible and use the solidus (/) instead of a horizontal line for small fractional terms, e.g., X/Y. In principle, variables are to be presented in italics. Powers of e are often more conveniently denoted by exp. Number consecutively any equations that have to be displayed separately from the text (if referred to explicitly in the text).

Footnotes

Footnotes should be used sparingly. Number them consecutively throughout the article. Many word processors can build footnotes into the text, and this feature may be used. Otherwise, please indicate the position of footnotes in the text and list the footnotes themselves separately at the end of the article. Do not include footnotes in the Reference list.

Artwork

Electronic Artwork

General points

- Make sure you use uniform lettering and sizing of your original artwork.
- Embed the used fonts if the application provides that option.
- Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately in the Figure Legends section of the manuscript..
- Size the illustrations close to the desired dimensions of the printed version.
- Submit each illustration as a separate file.

A detailed guide on electronic artwork is available on our website:

<http://www.elsevier.com/artworkinstructions>

You are urged to visit this site; some excerpts from the detailed information are given here.

Formats

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.

Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings, embed all used fonts.

TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.

TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.

TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

Please do not:

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

Color artwork

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color online (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or online only. [Further information on the preparation of electronic artwork.](#)

Illustration services

[Elsevier's WebShop](#) offers Illustration Services to authors preparing to submit a manuscript but concerned about the quality of the images accompanying their article. Elsevier's expert illustrators can produce scientific, technical and medical-style images, as well as a full range of charts, tables and graphs. Image 'polishing' is also available, where our illustrators take your image(s) and improve them to a professional standard. Please visit the website to find out more.

Figure legends

Ensure that each illustration has a legend. Supply legends separately, not attached to the figure. A legend should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

Tables

Please submit tables as editable text and not as images. Tables can be placed either next to the relevant text in the article, or on separate page(s) at the end. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules and shading in table cells.

Permissions

If any illustration or table has been previously published, a copy of the letter of PERMISSION to reprint from the copyright holder must accompany the manuscript (a sample Permission Request Form can be found at <http://www.elsevier.com/journal-authors/obtain-permission>). The source of the illustration or table should be included in the reference section of the manuscript. The legend should conclude with "Reprinted with permission" followed by the reference number, eg, "Reprinted with permission.23" Authors are responsible for any permission fees requested by the copyright holder.

Do not submit the same illustration or table simultaneously to two publishers on the assumption that one article will publish before the other and that the publisher of the first article will grant permission for reuse of the material. Publication of one article may be delayed unexpectedly, necessitating withdrawal of material from the other article at a late stage of production, which would delay publication of the issue.

References

Citation in text

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

Reference numbers (citations) in the text follow numerical order and should be superscripted. References are listed according to their order in the text, not in alphabetical order. The maximum number of references is 100.

Reference links

Increased discoverability of research and high quality peer review are ensured by online links to the sources cited. In order to allow us to create links to abstracting and indexing services, such as Scopus, CrossRef and PubMed, please ensure that data provided in the references are correct. Please note that incorrect surnames, journal/book titles, publication year and pagination may prevent link creation. When copying references, please be careful as they may already contain errors. Use of the DOI is encouraged.

A DOI can be used to cite and link to electronic articles where an article is in-press and full citation details are not yet known, but the article is available online. A DOI is guaranteed never to change, so you can use it as a permanent link to any electronic article. An example of a citation using DOI for an article not yet in an issue is: VanDecar J.C., Russo R.M., James D.E., Ambeh W.B., Franke M. (2003). Aseismic continuation of the Lesser Antilles slab beneath northeastern Venezuela. *Journal of Geophysical Research*, <https://doi.org/10.1029/2001JB000884>. Please note the format of such citations should be in the same style as all other references in the paper.

Please note that manuscripts in press may be referenced; however, manuscripts that have been submitted for publication but not yet accepted should not be referenced. All references must be complete when the manuscript is submitted.

Web references

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

Data references

This journal encourages you to cite underlying or relevant datasets in your manuscript by citing them in your text and including a data reference in your Reference List. Data references should include the following elements: author name(s), dataset title, data repository, version (where available), year, and global persistent identifier. Add [dataset] immediately before the reference so we can properly identify it as a data reference. The [dataset] identifier will not appear in your published article.

References in this issue

When referring to another article in the Seminars issue, incorporate it as an In Press article and the page numbers will be added by the editorial staff prior to printing.

Reference style

Text: Indicate references by (consecutive) superscript arabic numerals in the order in which they appear in the text. The numerals are to be used *outside* periods and commas, *inside* colons and semicolons. For further detail and examples you are referred to the [AMA Manual of Style](#), A Guide for Authors and Editors, Tenth Edition, ISBN 0-978-0-19-517633-9.

List: Number the references in the list in the order in which they appear in the text.

Examples:

Reference to a journal publication:

1. Van der Geer J, Hanraads JAJ, Lupton RA. The art of writing a scientific article. *J Sci Commun*. 2010;163:51–59.

Reference to a book:

2. Strunk W Jr, White EB. *The Elements of Style*. 4th ed. New York, NY: Longman; 2000.

Reference to a chapter in an edited book:

3. Mettam GR, Adams LB. How to prepare an electronic version of your article. In: Jones BS, Smith RZ, eds. *Introduction to the Electronic Age*. New York, NY: E-Publishing Inc; 2009:281–304.

Reference to a website:

4. Cancer Research UK. Cancer statistics reports for the UK. <http://www.cancerresearchuk.org/aboutcancer/statistics/cancerstatsreport/>; 2003 Accessed 13 March 2003.

Reference to a dataset:

[dataset] 5. Oguro, M, Imahiro, S, Saito, S, Nakashizuka, T. Mortality data for Japanese oak wilt disease and surrounding forest compositions, Mendeley Data, v1; 2015. <https://doi.org/10.17632/xwj98nb39r.1>.

Journal abbreviations source

Journal names should be abbreviated according to the [List of Title Word Abbreviations](#).

Video

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. . In order to ensure that your video or animation material is directly usable, please provide the file in one of our recommended file formats with a preferred maximum size of 150 MB per file, 1 GB in total. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including [ScienceDirect](#). Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our [video instruction pages](#). Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

Supplementary material

Supplementary material such as applications, images and sound clips, can be published with your article to enhance it. Submitted supplementary items are published exactly as they are received (Excel or PowerPoint files will appear as such online). Please submit your material together with the article and supply a concise, descriptive caption for each supplementary file. If you wish to make changes to supplementary material during any stage of the process, please make sure to provide an updated file. Do not annotate any corrections on a previous version. Please switch off the 'Track Changes' option in Microsoft Office files as these will appear in the published version.

Submission Checklist

The following list will be useful during the final checking of an article prior to sending it to the journal for review. Please consult this Guide for Authors for further details of any item.

• Ensure that the following items are present:

- One author has been designated as the corresponding author, with contact details:
- E-mail address
- Full postal address
- Phone numbers
- Disk is enclosed
- The electronic version and the hardcopy of the manuscript are identical
- Disk has been labelled with

article details (first author, first words of title)

file name(s)

media format (e.g., PC, Mac)

file format (e.g., Word, LaTeX)

- All text pages
- Keywords
- Original artwork (high-quality prints)
- All figure captions
- All tables (including title, description, footnotes)

Further considerations

- Manuscript has been 'spell-checked' and 'grammar-checked'
- References are in the correct format for this journal
- All references mentioned in the Reference list are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources (including the Web)
- Color figures are clearly marked as being intended for color reproduction on the Web (free of charge) and in print (a charge will apply), or to be reproduced in color on the Web (free of charge) and in black-and-white in print
- If only color on the Web is required, black and white versions of the figures are also supplied for printing purposes

For any further information please visit our customer support site at: <http://support.elsevier.com>.

AFTER ACCEPTANCE

Proofs

One set of page proofs (as PDF files) will be sent by e-mail to the corresponding author (if we do not have an e-mail address then paper proofs will be sent by post) or, a link will be provided in the e-mail so that authors can download the files themselves. Elsevier now provides authors with PDF proofs which can be annotated; for this you will need to [download the free Adobe Reader](#), version 9 (or higher). Instructions on how to annotate PDF files will accompany the proofs (also given online). The exact system requirements are given at the [Adobe site](#).

If you do not wish to use the PDF annotations function, you may list the corrections (including replies to the Query Form) and return them to Elsevier in an e-mail. Please list your corrections quoting line number. If, for any reason, this is not possible, then mark the corrections and any other comments (including replies to the Query Form) on a printout of your proof and scan the pages and return via e-mail. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. We will do everything possible to get your article published quickly and accurately. It is important to ensure that all corrections are sent back to us in one communication: please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

Three free copies of the issue will also be sent to the corresponding author of each article.

Offprints

The corresponding author will, at no cost, receive a customized [Share Link](#) providing 50 days free access to the final published version of the article on [ScienceDirect](#). The Share Link can be used for sharing the article via any communication channel, including email and social media. For an extra charge, paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any

time via Elsevier's [Webshop](#). Corresponding authors who have published their article open access do not receive a Share Link as their final published version of the article is available open access on ScienceDirect and can be shared through the article DOI link.

AUTHOR INQUIRIES

If you have any questions about the content of your article, its length, or the deadline, please contact the Guest Editor who has invited you to contribute.

If you have specific questions about manuscript preparation or publication procedures, contact: Jennifer Moulder, Journal Manager, Tel/Fax: (610) 321-9865, Email: j.moulder@elsevier.com

© Copyright 2018 Elsevier | <https://www.elsevier.com>